arXiv:q-alg/9703013v1 6 Mar 1997

q-alg/9703013

Factorisation of Macdonald polynomials [1]

V.B. Kuznetsov [†2] and E.K. Sklyanin [‡3]

[†] Department of Applied Mathematical Studies, University of Leeds, Leeds LS2 9JT, UK
[‡] Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606, Japan

1. Macdonald polynomials

Macdonald polynomials $P_\lambda(x;q,t)$ are orthogonal symmetric polynomials which are the natural multivariable generalisation of the *continuous q-ultraspherical polynomials* $C_n(x;\beta|q)$ [2] which, in their turn, constitute an important class of hypergeometric orthogonal polynomials in one variable. Polynomials $C_n(x;\beta|q)$ can be obtained from the general *Askey-Wilson polynomials* [3] through a specification of their four parameters (see, for instance, [9]), so that $C_n(x;\beta|q)$ depend only on one parameter β, apart from the degree n and the basic parameter q. In an analogous way, the Macdonald polynomials $P_\lambda(x;q,t)$ with one parameter t could be obtained as a limiting case of the 5-parameter Koornwinder's multivariable generalisation of the Askey-Wilson polynomials [10].

The main reference for the Macdonald polynomials is the book [17], Ch. VI, where they are called *symmetric functions with two parameters*. Let $\mathbb{K}=\mathbb{Q}(q,t)$ be the field of rational functions in two indeterminants q,t; $\mathbb{K}[x]=\mathbb{K}[x_1,\dots,x_n]$ be the ring of polynomials in n variables $x=(x_1,\dots,x_n)$ with coefficients in $\mathbb{K}$; and $\mathbb{K}[x]^W$ be the subring of all symmetric polynomials. The Macdonald polynomials $P_\lambda(x)=P_\lambda(x;q,t)$ are symmetric polynomials labelled by the sequences $\lambda=\{0\le\lambda_1\le\lambda_2\le\ldots\le\lambda_n\}$ of integers (dominant weights). They form a $\mathbb{K}$ basis of $\mathbb{K}[x]^W$ and are uniquely characterised as joint eigenvectors of the commuting q-difference operators H_k

$$H_k\,P_\lambda = h_{k;\lambda}\,P_\lambda\,,\qquad k=1,\dots,n\,, \tag{1.1}$$

normalised by the condition

$$P_\lambda=\sum_{\lambda'\preceq\lambda}\kappa_{\lambda\lambda'}\,m_{\lambda'},\qquad \kappa_{\lambda\lambda}=1\,,\qquad (\kappa_{\lambda\lambda'}\in\mathbb{K}) \tag{1.2}$$

[1] Talk given on July 4th, 1996 on the Second Workshop "Symmetries and Integrability of Difference Equations", Canterbury, UK.

[2] On leave from: Department of Mathematical and Computational Physics, Institute of Physics, St. Petersburg University, St. Petersburg 198904, Russian Federation. E-mail: vadim@amsta.leeds.ac.uk

[3] On leave from: Steklov Mathematical Institute, Fontanka 27, St. Petersburg 191011, Russian Federation. E-mails: sklyanin@kurims.kyoto-u.ac.jp and sklyanin@pdmi.ras.ru

where, for each μ, $m_\mu(x)$ stands for the monomial symmetric function: $m_\mu(x) = \sum x_1^{\nu_1} \dots x_n^{\nu_n}$ with the sum taken over all distinct permutations ν of μ, and $\preceq$ is the dominance order on the dominant weights

$$\lambda' \preceq \lambda \quad \Longleftrightarrow \quad \left\{ |\lambda'| = |\lambda| \; ; \quad \sum_{j=k}^{n} \lambda'_j \le \sum_{j=k}^{n} \lambda_j \, , \quad k = 2, \dots, n \right\} .$$

The commuting operators H_i have the form:

$$H_i = \sum_{\substack{J \subset \{1,\dots,n\} \\ |J|=i}} \left(\prod_{\substack{j \in J \\ k \in \{1,\dots,n\} \setminus J}} v_{jk} \right) \left(\prod_{j \in J} T_{q,x_j} \right) , \qquad i = 1, \dots, n \, , \tag{1.3}$$

where

$$v_{jk} = \frac{t^{\frac{1}{2}} x_j - t^{-\frac{1}{2}} x_k}{x_j - x_k} \, , \qquad |q| < 1 \, , \; |t| < 1 \, . \tag{1.4}$$

The T_{q,x_j} stands for the q-shift operator in the variable x_j: $(T_{q,x_j} f)(x_1, \dots, x_n) = f(x_1, \dots, \, qx_j, \, \dots, x_n)$. The operators (1.3) were introduced for the first time in [21] and are the integrals of motion for the quantum Ruijsenaars model, which is a relativistic (or q-) analog of the trigonometric Calogero-Moser-Sutherland model. They are called sometimes Macdonald operators in the mathematical literature.

The corresponding eigenvalues $h_{k;\lambda}$ in (1.1) are

$$h_{k;\lambda} = \sum_{j_1 < \dots < j_k} \mu_{j_1} \dots \mu_{j_k} \, , \qquad \mu_j = q^{\lambda_j} t^{j - \frac{n+1}{2}} \, . \tag{1.5}$$

The polynomials P_λ are orthogonal

$$\frac{1}{(2\pi i)^n} \oint\limits_{|x_1|=1} \frac{dx_1}{x_1} \dots \oint\limits_{|x_n|=1} \frac{dx_n}{x_n} \, \overline{P_\lambda(x;q,t)} \, P_{\lambda'}(x;q,t) \, \Delta(x) = 0, \qquad \lambda \neq \lambda' \tag{1.6}$$

with respect to the weight

$$\Delta(x_1, \dots, x_n) = \prod_{j \neq k} \frac{(x_j x_k^{-1}; q)_\infty}{(t x_j x_k^{-1}; q)_\infty} \, . \tag{1.7}$$

For instance, for $n = 3$,

$$m_{000} = 1, \quad m_{001} = x_1 + x_2 + x_3, \quad m_{011} = x_1x_2 + x_1x_3 + x_2x_3, \quad m_{002} = x_1^2 + x_2^2 + x_3^2,$$

$$m_{111} = x_1x_2x_3, \quad m_{012} = x_1x_2^2 + x_1^2x_2 + x_1x_3^2 + x_1^2x_3 + x_2x_3^2 + x_2^2x_3,$$

$$m_{112} = x_1^2x_2x_3 + x_1x_2^2x_3 + x_1x_2x_3^2, \quad m_{022} = x_1^2x_2^2 + x_1^2x_3^2 + x_2^2x_3^2, \quad m_{003} = x_1^3 + x_2^3 + x_3^3.$$

$$P_{000} = m_{000}, \quad P_{001} = m_{001}, \quad P_{011} = m_{011}, \quad P_{002} = m_{002} + \tfrac{(1-t)(1+q)}{1-qt} \, m_{011},$$

$$P_{111} = m_{111}, \quad P_{012} = m_{012} + \tfrac{(1-t)(q(2t+1)+t+2)}{1-qt^2} \, m_{111},$$

$$P_{112} = m_{112}, \quad P_{022} = m_{022} + \tfrac{(1-t)(1+q)}{1-qt} \, m_{112},$$

$$P_{003} = m_{003} + \tfrac{(1-t)(1+q+q^2)}{1-q^2t}\, m_{012} + \tfrac{(1-t)^2(1+q)(1+q+q^2)}{(1-qt)(1-q^2t)}\, m_{111}.$$

Rodrigues type of formula for Macdonald polynomials was recently found in [11, 12, 16]. In the limit $q \uparrow 1$ Macdonald polynomials turn into *Jack polynomials* [17].

A multivariable function/polynomial can be called *special function* if it is some recognised (classical) special function in the case of one variable and if it is common eigenfunction of a complete set of commuting linear partial differential/difference operators defining a *quantum integrable system*. Macdonald polynomials *are* special functions in the above-mentioned sense. They diagonalise the integrals of motion H_j of the quantum Ruijsenaars system.

For any special function in many variables one can set up a general problem of its factorisation in terms of one-variable (special) functions. For the Macdonald polynomials P_λ this would mean finding a factorising integral operator M such that

$$M : P_\lambda(x; q, t) \mapsto \prod_{j=1}^{n} \Phi_{\lambda,j}(y_j)\,.$$

For some particular special functions (or, in other words, for some particular quantum integrable systems) such an operator might simplify to a local transform which is a simple change of variables, from x to y. This happens for example in the case of ellipsoidal harmonics in $\mathbb{R}^n$ (see [18] and [23] for $n = 3$), correspondingly, in the case of quantum Neumann system, where the transform $x \mapsto y$ is the change of variables, from Cartesian to ellipsoidal.

As it is shown further on, in Section 3, the factorising operator for the (symmetric) Macdonald polynomials has to be non-local, i.e. to be some linear integral operator. Moreover, it is explicitly described in the first two non-trivial cases (when $n = 2$ and $n = 3$) in terms of the Askey-Wilson operator.

The formulated factorisation problem also makes sense in the limit to the Liouville integrable systems in classical mechanics, becoming there the well-known problem of separation of variables (SoV) in the Hamilton-Jacobi equation.

The kernel of the factorising integral operator M turns in this limit into the generating function of the separating canonical transformation.

2. Hypergeometric polynomials f_λ and φ_λ in one variable

In this Section we introduce the hypergeometric polynomials $f_\lambda/\varphi_\lambda$ of one variable each constituting a basis which is conjugated to Jack/Macdonald polynomials with respect to the factorising integral transform M. First of all, we describe the procedure of lowering the order of (basic) hypergeometric functions which will lead us to these new sets of interesting polynomials in one variable. We will use (becoming already) standard notations of [8] for the (basic) hypergeometric series and other formulas of the q-analysis.

In 1927 Fox [6] found an interesting relation between hypergeometric functions:

$$ {}_pF_q\left[\begin{matrix} a_1+m_1, a_2, \dots, a_p \\ b_1, \dots, b_q \end{matrix}; y\right] = \sum_{j=0}^{\infty} \frac{(-y)^j (-m_1)_j}{j!} \frac{(a_2)_j \cdots (a_p)_j}{(b_1)_j \cdots (b_q)_j} \; {}_pF_q\left[\begin{matrix} a_1+j, \dots, a_p+j \\ b_1+j, \dots, b_q+j \end{matrix}; y\right] . \tag{2.1} $$

When $a_1 = b_1$ (2.1) gives the expansion of ${}_pF_q\left[\begin{matrix} a \\ b \end{matrix}; y\right]$ in terms of functions of the type ${}_{p-1}F_{q-1}\left[\begin{matrix} a \\ b \end{matrix}; y\right]$. When m_1 is a positive integer, the series on the right of (2.1) terminates, and we have (for the case of $a_1 = b_1$ and $q = p-1$) the following relation

$$ {}_pF_{p-1}\left[\begin{matrix} b_1+m_1, a_2, \dots, a_p \\ b_1, b_2 \dots, b_{p-1} \end{matrix}; y\right] \tag{2.2} $$

$$ = \sum_{j=0}^{m_1} \frac{(-y)^j (-m_1)_j}{j!} \frac{(a_2)_j \cdots (a_p)_j}{(b_1)_j \cdots (b_{p-1})_j} \; {}_{p-1}F_{p-2}\left[\begin{matrix} a_2+j, \dots, a_p+j \\ b_2+j, \dots, b_{p-1}+j \end{matrix}; y\right] . $$

Supposing that $a_2 = b_2 + m_2$, $a_3 = b_3 + m_3, \dots,$ $a_{p-1} = b_{p-1} + m_{p-1}$ with some non-negative integers m_k, $k = 2, \dots, p-1$, we then iterate the relation (2.2) further on and, finally, using the binomial theorem,

$$ {}_1F_0\left[\begin{matrix} a \\ - \end{matrix}; y\right] = (1-y)^{-a} , $$

conclude that the function

$$ f_{m_1,\dots,m_{p-1}} \equiv (1-y)^{a_p + \sum_{j=1}^{p-1} m_j} \; {}_pF_{p-1}\left[\begin{matrix} b_1+m_1, b_2+m_2, \dots, b_{p-1}+m_{p-1}, a_p \\ b_1, b_2 \dots, b_{p-1} \end{matrix}; y\right] $$

is a polynomial in y of the cumulative degree $\sum_{j=1}^{n-1} m_j$.

As for a q-analog of the reduction formula (2.2) we refer to (1.9.4) in [8] (see also [7]). It was proved for the first time in [4] in a more general case (although in different notations than ones in [8] which are adopted here). Following [8], consider the q-analog of the Vandermonde formula in the form

$$ {}_2\phi_1\left[\begin{matrix} q^{-n}, q^{-m} \\ b_{p-1} \end{matrix}; q, q\right] = \frac{(b_{p-1}q^m; q)_n}{(b_{p-1}; q)_n} \; q^{-mn} \tag{2.3} $$

where m and n are non-negative integers such that $m \geq n$, then we can reduce the order of the basic hypergeometric function through the following equalities ($|y| < 1$):

$$ {}_p\phi_{p-1}\left[\begin{matrix} a_1, \dots, a_{p-1}, b_{p-1}q^m \\ b_1, \dots, b_{p-2}, b_{p-1} \end{matrix}; q, y\right] \tag{2.4} $$

$$ = \sum_{n=0}^{\infty} \frac{(a_1,\dots,a_{p-1};q)_n}{(q,b_1,\dots,b_{p-2};q)_n} \; y^n \sum_{k=0}^{n} \frac{(q^{-n},q^{-m};q)_k}{(q,b_{p-1};q)_k} \; q^{mn+k} $$

$$ = \sum_{n=0}^{\infty} \sum_{k=0}^{m} \frac{(a_1,\dots,a_{p-1};q)_n (q^{-m};q)_k}{(b_1,\dots,b_{p-2};q)_n (q;q)_{n-k} (q,b_{p-1};q)_k} \; y^n \; (-1)^k q^{mn+k-nk+\binom{k}{2}} $$

$$ = \sum_{k=0}^{m} \frac{(q^{-m},a_1,\dots,a_{p-1};q)_k}{(q,b_1,\dots,b_{p-1};q)_k} \; (-yq^m)^k q^{-\binom{k}{2}} \; {}_{p-1}\phi_{p-2}\left[\begin{matrix} a_1q^k, \dots, a_{p-1}q^k \\ b_1q^k, \dots, b_{p-2}q^k \end{matrix}; q, yq^{m-k}\right] . $$

Then again, iterating it and using the q-binomial theorem,

$$ {}_1\phi_0\left[\begin{matrix} a \\ - \end{matrix}; q, y\right] = \frac{(ay;q)_\infty}{(y;q)_\infty}\,, \qquad |y|<1\,, \qquad |q|<1\,, $$

we conclude that, for non-negative integers m_k, $k=1,\dots,p-1$, the function

$$ \varphi_{m_1,\dots,m_{p-1}} \equiv \frac{(y;q)_\infty}{(ya_pq^{m_1+\dots+m_{p-1}};q)_\infty}\ {}_p\phi_{p-1}\left[\begin{matrix} b_1q^{m_1},\dots,b_{p-1}q^{m_{p-1}},a_p \\ b_1,\dots,b_{p-1} \end{matrix}; q, y\right] $$

is a polynomial in y of the cumulative degree $\sum_{j=1}^{n-1} m_j$.

Now we can connect integers m_j to a partition $\lambda=\{\lambda_1,\dots,\lambda_n\}$, namely:

$$ m_j = \lambda_{j+1}-\lambda_j \quad (\geq 0)\,, \qquad j=1,\dots,n-1\,. $$

Put also ($g\in\mathbb{R}$)

$$ b_j = \lambda_1-\lambda_{j+1}+1-jg\,, \quad a_j=b_j+m_j\,, \quad j=1,\dots,n-1\,, \quad a_n=\lambda_1-\lambda_n+1-ng\,. $$

Let us define the following function

$$ f_\lambda(y) := y^{\lambda_1}\,(1-y)^{1-ng}\ {}_nF_{n-1}\left[\begin{matrix} b_1+m_1,\dots,b_{n-1}+m_{n-1},a_n \\ b_1,\dots,b_{n-1} \end{matrix}; y\right]\,. $$

Obviously, this function is a polynomial in y of the form

$$ \sum_{k=\lambda_1}^{\lambda_n} \chi_k\, y^k\,. \tag{2.5} $$

The polynomials $f_\lambda(y)$ were introduced in [13] and they are factorised polynomials of one variable for the multivariable Jack polynomials. In [22, 13] the corresponding sine-kernel (of Gegenbauer type) factorising the A_2 Jack polynomials was described in detail. In the sequel we concentrate on the q-analogs of these polynomials, the $\varphi_\lambda(y)$, which were introduced in [14].

Put again $m_j=\lambda_{j+1}-\lambda_j$, $j=1,\dots,n-1$, but

$$ b_j = q^{\lambda_1-\lambda_{j+1}+1-jg}\,, \quad a_j=b_jq^{m_j}\,, \quad j=1,\dots,n-1\,, \quad a_n=q^{\lambda_1-\lambda_n+1-ng}\,. $$

We will also use the parameter t which is connected to g in the following way

$$ t := q^g\,. $$

Then the polynomials φ_λ are defined as follows:

$$ \varphi_\lambda(y) := y^{\lambda_1}\,(y;q)_{1-ng}\ {}_n\phi_{n-1}\left[\begin{matrix} b_1q^{m_1},\dots,b_{n-1}q^{m_{n-1}},a_n \\ b_1,\dots,b_{n-1} \end{matrix}; q, y\right]\,, \tag{2.6} $$

and, like $f_\lambda(y)$, expand in y as (2.5).

In [14] we have found several useful representations for these polynomials in addition to their definition (2.6). Let us list here some of them. Introduce notations:

$$\lambda_{ij} := \lambda_i - \lambda_j\,, \qquad |\lambda| = \sum_{j=1}^{n} \lambda_j\,.$$

First of all, the coefficients χ_k in (2.5) have the following explicit representation:

$$\chi_k = \left(q^{-1}t^n\right)^{\lambda_1-k} \frac{(q^{-1}t^n;q)_{k-\lambda_1}}{(q;q)_{k-\lambda_1}}\; {}_{n+1}\phi_n\left[\begin{matrix} q^{\lambda_1-k}, a_1, \ldots, a_n \\ q^{\lambda_1-k+2}t^{-n}, b_1, \ldots, b_{n-1}\end{matrix}; q, q\right]. \tag{2.7}$$

It is easy to give simpler expressions for some of χ_k such as

$$\chi_{\lambda_1} = 1, \qquad \chi_{\lambda_n} = t^{n\lambda_1-|\lambda|} \prod_{j=1}^{n-1} \frac{(t^j;q)_{\lambda_j-\lambda_1}(t^j;q)_{\lambda_n-\lambda_{n-j}}}{(t^j;q)_{\lambda_{j+1}-\lambda_1}(t^j;q)_{\lambda_n-\lambda_{n-j+1}}}\,. \tag{2.8}$$

Let us give few first polynomials in the case $n = 3$

$$\varphi_{000} = 1, \quad \varphi_{001} = 1 + \tfrac{1}{t(1+t)}\, y, \quad \varphi_{011} = 1 + \tfrac{1+t}{t^2}\, y\,,$$

$$\varphi_{002} = 1 + \tfrac{(1+q)(1-t)}{t(1-qt^2)}\, y + \tfrac{1-qt}{t^2(1-qt^2)(1+t)}\, y^2\,,$$

$$\varphi_{012} = 1 + \tfrac{1+qt+t-qt^2-t^2-qt^3}{t^2(1-qt^2)}\, y + \tfrac{1}{t^3}\, y^2\,,$$

$$\varphi_{022} = 1 + \tfrac{(1+q)(1-t^2)}{t^2(1-qt)}\, y + \tfrac{(1+t)(1-qt^2)}{t^4(1-qt)}\, y^2\,,$$

$$\varphi_{003} = 1 + \tfrac{(1+q+q^2)(1-t)}{t(1-q^2t^2)}\, y + \tfrac{(1+q+q^2)(1-t)}{t^2(1+qt)(1-qt^2)}\, y^2 + \tfrac{1-q^2t}{t^3(1+t)(1+qt)(1-qt^2)}\, y^3\,.$$

There is also a simple formula for $\varphi_\lambda(t^n)$

$$\varphi_\lambda(t^n) = t^{n\lambda_1}(t^n;q)_{\lambda_{n1}} \prod_{j=1}^{n-1} \frac{(t^j;q)_{\lambda_j-\lambda_1}}{(t^j;q)_{\lambda_{j+1}-\lambda_1}}\,. \tag{2.9}$$

There is a nice representation of the polynomials φ_λ in terms of the ϕ_D q-Lauricella function [1, 19, 5]

$$\varphi_\lambda(y) = y^{\lambda_1}\, \frac{(qt^{-n}q^{\lambda_{1n}}y;q)_{\lambda_{n1}}}{\prod_{j=1}^{n-1}(q^{\lambda_1-\lambda_{n-j+1}+1}t^{j-n};q)_{\lambda_{n-j+1}-\lambda_{n-j}}}\; \phi_D\left[\begin{matrix} y; b'_1, \ldots, b'_{n-1} \\ qt^{-n}q^{\lambda_{1n}}y\end{matrix}; q; a'_1, \ldots, a'_{n-1}\right],$$

with

$$a'_j = qt^{j-n}q^{\lambda_1-\lambda_{n-j}}, \qquad b'_j = q^{\lambda_{n-j}-\lambda_{n-j+1}}\,, \qquad j = 1, \ldots, n-1\,. \tag{2.10}$$

The ϕ_D q-Lauricella function of $n-1$ variables z_i is a multivariable generalisation of the basic hypergeometric series and is defined by the formula

$$\phi_D\left[\begin{matrix} a'; b'_1, \ldots, b'_{n-1} \\ c\end{matrix}; q; z_1, \ldots, z_{n-1}\right] := \sum_{k_1,\ldots,k_{n-1}=0}^{\infty} \frac{(a';q)_{k_1+\ldots+k_{n-1}}}{(c;q)_{k_1+\ldots+k_{n-1}}} \prod_{j=1}^{n-1} \frac{(b'_j;q)_{k_j} z_j^{k_j}}{(q;q)_{k_j}}\,,$$

using which we get the most explicit representation of our polynomials φ_λ

$$\varphi_\lambda(y) = y^{\lambda_1} \left(\prod_{j=1}^{n-1} (q^{\lambda_1-\lambda_{n-j+1}+1} t^{j-n}; q)_{\lambda_{n-j+1}-\lambda_{n-j}} \right)^{-1} \times$$

$$\times \sum_{k_1=0}^{\lambda_n-\lambda_{n-1}} \cdots \sum_{k_{n-1}=0}^{\lambda_2-\lambda_1} (qt^{-n} q^{\lambda_{1n}+k_1+\ldots+k_{n-1}} y; q)_{\lambda_{n1}-k_1-\ldots-k_{n-1}} \, (y;q)_{k_1+\ldots+k_{n-1}}$$

$$\times \prod_{j=1}^{n-1} \frac{(q^{\lambda_{n-j}-\lambda_{n-j+1}}; q)_{k_j} (qt^{j-n} q^{\lambda_1-\lambda_{n-j}})^{k_j}}{(q;q)_{k_j}} \,. \tag{2.11}$$

Finally, these polynomials satisfy the following q-difference equation

$$\sum_{k=0}^{n} (-1)^k \, t^{-\frac{n-1}{2}k} \, (1 - q^k t^{-k} y) \, (y;q)_k \, (q^{k+1} t^{-n} y; q)_{n-k} \, h_{n-k;\lambda} \, \varphi_\lambda(q^k y) = 0 \tag{2.12}$$

where $h_{k;\lambda}$ are given by (1.5) and we assume $h_{0;\lambda} \equiv 1$.

Let us give few remarks about this new class of basic hypergeometric polynomials in one variable.

Remark 1. First of all, we stress that our way of extracting polynomials from hypergeometric series is quite different from the usual one. Indeed, all classical orthogonal q-polynomials of one variable are obtained by just terminating the corresponding hypergeometric series. For instance, the generic Askey-Wilson orthogonal q-polynomials which appear on the ${}_4\phi_3$ level are defined as follows

$$p_n(x; a, b, c, d|q) = const \cdot {}_4\phi_3 \left[\begin{matrix} q^{-n}, abcdq^{n-1}, ae^{i\theta}, ae^{-i\theta} \\ ab, ac, ad \end{matrix} ; q, q \right] , \qquad x = \cos\theta \,.$$

In contrast, to extract the polynomials φ_λ at the level ${}_n\phi_{n-1}$ we use the procedure of order reduction (2.4) of the basic hypergeometic functions with the specific choice of upper and lower parameters, namely: when the ratio of an upper parameter and one of the lower parameters is equal to q^{m_i} where m_i are non-negative integers. The procedure of order reduction is thus an important second possibility (in addition to simple termination) in order to get polynomials out of hypergeometric series.

Remark 2. The importance of polynomials φ_λ stems from the fact that they are the *factorised polynomials* of one variable for the multivariable Macdonald polynomials. Notice that the polynomials φ_λ, as well as Macdonald polynomials P_λ, are labelled by the dominant weights. The multivariable polynomials Φ_λ combined from the one-variable polynomials φ_λ:

$$\Phi_\lambda(y_1, \ldots, y_n) := y_n^{|\lambda|} \prod_{k=1}^{n-1} \varphi_\lambda(y_k) \,,$$

satisfy the following multiparameter spectral problem (cf. (2.12))

$$\Phi_\lambda(y_1, \ldots, y_{n-1}, qy_n) = h_{n;\lambda} \, \Phi_\lambda(y_1, \ldots, y_n) \,, \tag{2.13}$$

$$\sum_{k=0}^{n} (-1)^k\, t^{-\frac{n-1}{2}k}\, (1-q^k t^{-k} y_j)\, (y_j;q)_k\, (q^{k+1}t^{-n}y_j;q)_{n-k}\, h_{n-k;\lambda} \times$$

$$\times \Phi_\lambda(y_1,\ldots,q^k y_j,\ldots,y_n) = 0\,, \qquad j=1,\ldots,n-1\,,$$

with *the same set of spectral parameters* $(h_{1;\lambda},\ldots,h_{n;\lambda})$ (1.5) as in the spectral problem (1.1) for the Macdonald polynomials P_λ. Hence, one can introduce the commuting operators $H_i^{(y)}$, $i=1,\ldots,n$ defined by their eigenfunctions $\Phi_\lambda(y)$ and eigenvalues $h_{k;\lambda}$. Since the spectrum $(h_{1;\lambda},\ldots,h_{n;\lambda})$ coincides with that (1.5) of the Macdonald polynomials, the two sets of commuting operators are isomorphic and there has to exist an intertwining linear operator M

$$M\; H_i^{(x)} = H_i^{(y)}\; M \qquad (H_i^{(x)} \equiv H_i)\,.$$

Actually, to any choice of the normalisation coefficients c_λ in

$$M:\; P_\lambda(x) \mapsto c_\lambda\; \Phi_\lambda(y)$$

there corresponds some intertwiner M. The problem is to select a factorising operator M having an explicit description in terms of its integral kernel or its matrix in some basis in $\mathbb{K}[x]^W$. In [14, 15] we have found such an operator as well as its inversion in the first two non-trivial cases, when $n=2$ and $n=3$. It appears that the factorising operator M can be expressed in those cases through the Askey-Wilson operator.

3. The cases $n=2$ and $n=3$

Let us first describe the integral operator M_ξ performing the separation of variables in the A_1 Macdonald polynomials (we skip the trivial case of the purely coordinate SoV $x_{1,2} \to x_\pm \equiv (x_1 x_2^{\pm 1})^{1/2}$). Our main technical tool is the famous Askey-Wilson integral identity [3, 8]

$$\frac{1}{2\pi i} \int\limits_{\Gamma_{abcd}} \frac{dx}{x}\, \frac{(x^2,x^{-2};q)_\infty}{(ax,ax^{-1},bx,bx^{-1},cx,cx^{-1},dx,dx^{-1};q)_\infty} = \frac{2(abcd;q)_\infty}{(q,ab,ac,ad,bc,bd,cd;q)_\infty}\,. \tag{3.1}$$

The cycle Γ_{abcd} depends on complex parameters a,b,c,d and is defined as follows. Let $C_{z,r}$ be the counter-clockwise oriented circle with the center z and radius r. If $\max(|a|\,,|b|\,,|c|\,,\,|d|\,,|q|) < 1$ then $\Gamma_{abcd} = C_{0,1}$. The identity (3.1) can be continued analytically for the values of parameters a, b, c, d outside the unit circle provided the cycle Γ_{abcd} is deformed appropriately. In general case

$$\Gamma_{abcd} = C_{0,1} + \sum_{z=a,b,c,d} \sum_{\substack{k\geq 0 \\ |z|q^k>1}} (C_{zq^k,\varepsilon} - C_{z^{-1}q^{-k},\varepsilon})\,,$$

ε being small enough for $C_{z^{\pm 1}q^{\pm k},\varepsilon}$ to encircle only one pole of the denominator.

Put

$$a = yq^{\frac{\alpha}{2}}\,, \qquad b = y^{-1}q^{\frac{\alpha}{2}}\,, \qquad c = rq^{\frac{\beta}{2}}\,, \qquad d = r^{-1}q^{\frac{\beta}{2}}\,.$$

We will use the notation $\Gamma^{ry}_{\alpha\beta}$ for the contour obtained from Γ_{abcd} by these substitutuions. Introduce also the following useful notation

$$\mathcal{L}_q(\nu; x, y) := (\nu xy, \nu xy^{-1}, \nu x^{-1}y, \nu x^{-1}y^{-1}; q)_\infty\,. \tag{3.2}$$

Now let

$$\alpha = \beta = g\,, \quad y = y_-\,, \quad x = x_-\,, \quad r = t^{-1}y_+\,, \quad x_\pm \equiv (x_1 x_2^{\pm 1})^{\frac{1}{2}}\,, \quad y_\pm \equiv (y_1 y_2^{\pm 1})^{\frac{1}{2}}\,.$$

Introduce the kernel $\mathcal{M}(y_+, y_-|x_-)$:

$$\mathcal{M}(y_+, y_-|x_-) = \frac{(1-q)(q;q)_\infty^2\ (x_-^2, x_-^{-2}; q)_\infty\ \mathcal{L}_q(t; y_-, t^{-1}y_+)}{2B_q(g,g)\ \mathcal{L}_q\Big(t^{\frac{1}{2}}; y_-, x_-\Big)\ \mathcal{L}_q\Big(t^{\frac{1}{2}}; x_-, t^{-1}y_+\Big)}\,. \tag{3.3}$$

Assuming ξ to be an arbitrary complex parameter, we introduce the integral operator M_ξ acting on functions $f(x_1, x_2)$ by the formula

$$(M_\xi f)(y_1, y_2) \equiv \frac{1}{2\pi i} \int\limits_{\Gamma_{g,g}^{t^{-1}y_+, y_-}} \frac{dz}{z}\ \mathcal{M}(y_+, y_-|z)\ f(t^{-\frac{1}{2}}\xi y_+ z, t^{-\frac{1}{2}}\xi y_+ z^{-1})\,. \tag{3.4}$$

Theorem 1 ([15]) *The operator M_ξ (3.4),(3.3) performs the factorisation of (or, in other words, the SoV for) the A_1 Macdonald polynomials:*

$$M_\xi : P_\lambda(x_1, x_2) \to c_{\lambda,\xi}\ \varphi_\lambda(y_1)\ \varphi_\lambda(y_2)\,, \tag{3.5}$$

where $\varphi_\lambda(y)$ is the factorised (or separated) polynomial and the normalisation coefficient $c_{\lambda,\xi}$ is equal to

$$c_{\lambda,\xi} = t^{-2\lambda_1+\lambda_2}\xi^{|\lambda|}\ \frac{(t;q)_{\lambda_{21}}}{(t^2;q)_{\lambda_{21}}}\,. \tag{3.6}$$

Note that the relation (3.5) is equivalent [15] to the product formula for the continuous q-ultraspherical polynomials [20, 8].

The kernel of the inverse operator M_ξ^{-1} has the form

$$\widetilde{\mathcal{M}}(x_+, x_-|y_-) = \frac{(1-q)(q;q)_\infty^2\ (y_-^2, y_-^{-2}; q)_\infty\ \mathcal{L}_q\Big(t^{\frac{1}{2}}; x_-, t^{-\frac{1}{2}}\xi^{-1}x_+\Big)}{2B_q(-g, 2g)\ \mathcal{L}_q\Big(t^{-\frac{1}{2}}; y_-, x_-\Big)\ \mathcal{L}_q\Big(t; y_-, t^{-\frac{1}{2}}\xi^{-1}x_+\Big)}\,,$$

with the following substitutions for the contour Γ:

$$\alpha = -g\,, \qquad \beta = 2g\,, \qquad x = y_-\,, \qquad y = x_-\,, \qquad r = t^{-\frac{1}{2}}\xi^{-1}\,.$$

Theorem 2 ([15]) *The inversion of the operator M_ξ (3.4) is given by the formula*

$$(M_\xi^{-1} f)(x_1,x_2) = \frac{1}{2\pi i} \int\limits_{\Gamma_{-g,2g}^{t^{-1/2}\xi^{-1}x_+,x_-}} \frac{dz}{z}\, \widetilde{\mathcal{M}}(x_+,x_-|z)\; f(t^{\frac12}\xi^{-1}x_+z, t^{\frac12}\xi^{-1}x_+z^{-1})\,. \tag{3.7}$$

The operator M_ξ^{-1} provides an integral representation for the A_1 Macdonald polynomials in terms of the factorised polynomials $\varphi_{\lambda_1,\lambda_2}(y)$

$$M_\xi^{-1} : c_{\lambda,\xi}\; \varphi_\lambda(y_1)\; \varphi_\lambda(y_2) \to P_\lambda(x_1,x_2)\,. \tag{3.8}$$

In contrast to the formula (3.5) which paraphrases an already known result, the formula (3.8) leads to a new integral relation for the q-ultraspherical polynomials. Note that for positive integer g the operator M_ξ^{-1} becomes a q-difference operator of the order g (cf. [14] and [15]).

Now we describe the factorising operator M and its inversion in the case of A_2 Macdonald polynomials. Introduce the following operator M acting on functions $f(x_1,x_2,x_3)$ by the formula

$$\begin{aligned}(Mf)(y_1,y_2,y_3) \;&=\; \frac{1}{2\pi i} \int\limits_{\Gamma_{g,2g}^{t^{-\frac32}y_+,y_-}} \frac{dx_-}{x_-}\, \mathcal{M}((y_1y_2)^{\frac12},(y_1/y_2)^{\frac12}\,|\,x_-)\;\times \\ &\qquad \times f(t^{-\frac32}y_3(y_1y_2)^{\frac12}x_-, t^{-\frac32}y_3(y_1y_2)^{\frac12}x_-^{-1}, y_3)\end{aligned}$$

with the kernel

$$\mathcal{M}(y_+,y_-\mid x_-) = \frac{(1-q)(q;q)_\infty^2 (x_-^2,x_-^{-2};q)_\infty\, \mathcal{L}_q\Big(t^{\frac32};y_-,y_+t^{-\frac32}\Big)}{2B_q(g,2g)\,\mathcal{L}_q\Big(t^{\frac12};y_-,x_-\Big)\,\mathcal{L}_q\Big(t;x_-,y_+t^{-\frac32}\Big)}$$

and the following substitutions: $\alpha = g$, $\beta = 2g$, $r = t^{-\frac32}y_+$, $y = y_-$, $x = x_- \equiv (x_1x_2^{-1})^{\frac12}$, $x_+ \equiv (\frac{x_1x_2}{x_3^2})^{\frac12}$, $y_\pm \equiv (y_1y_2^{\pm1})^{\frac12}$.

Theorem 3 ([14]) *The operator M transforms any A_2 Macdonald polynomial P_λ into the product*

$$M : P_\lambda(x_1,x_2,x_3;q,t) \to c_\lambda\; y_3^{\lambda_1+\lambda_2+\lambda_3}\; \varphi_\lambda(y_1)\; \varphi_\lambda(y_2)$$

of factorised polynomials $\varphi_{\lambda_1\lambda_2\lambda_3}(y)$ of one variable

$$\varphi_\lambda(y) = y^{\lambda_1}\; (y;q)_{1-3g}\; {}_3\phi_2\left[\begin{matrix} t^{-3}q^{1-\lambda_{31}}, t^{-2}q^{1-\lambda_{21}}, t^{-1}q \\ t^{-2}q^{1-\lambda_{31}}, t^{-1}q^{1-\lambda_{21}}\end{matrix}\;;q,y\right].$$

The normalisation coefficient c_λ equals

$$c_\lambda = t^{\lambda_2-4\lambda_1}\; \frac{(t^2;q)_{\lambda_{31}}(t^2;q)_{\lambda_{32}}(t;q)_{\lambda_{21}}}{(t^3;q)_{\lambda_{31}}(t;q)_{\lambda_{32}}(t^2;q)_{\lambda_{21}}}\,.$$

Theorem 4 ([14]) *The inverse operator M^{-1} is the integral operator*

$$(M^{-1}f)(x_1,x_2,x_3)=\frac{1}{2\pi i}\int\limits_{\Gamma^{x_+,x_-}_{-g,3g}}\frac{dy_-}{y_-}\,\widetilde{\mathcal{M}}\left(\frac{(x_1x_2)^{\frac12}}{x_3},\left(\frac{x_1}{x_2}\right)^{\frac12}\,\Big|\,y_-\right)\times$$

$$\times f\left(\frac{t^{\frac32}(x_1x_2)^{\frac12}y_-}{x_3},\frac{t^{\frac32}(x_1x_2)^{\frac12}}{x_3y_-},x_3\right)$$

with the kernel

$$\widetilde{\mathcal{M}}(x_+,x_-\mid y_-)=\frac{(1-q)(q;q)_\infty^2(y_-^2,y_-^{-2};q)_\infty\,\mathcal{L}_q(t;x_-,x_+)}{2B_q(-g,3g)\,\mathcal{L}_q\Big(t^{-\frac12};y_-,x_-\Big)\,\mathcal{L}_q\Big(t^{\frac32};y_-,x_+\Big)}\,. \tag{3.9}$$

It provides a new integral representation for the A_2 Macdonald polynomials in terms of the factorised polynomials $\varphi_{\lambda_1,\lambda_2,\lambda_3}(y)$ of one variable

$$M^{-1}:\ c_\lambda\,y_3^{\lambda_1+\lambda_2+\lambda_3}\,\varphi_\lambda(y_1)\,\varphi_\lambda(y_2)\mapsto P_\lambda(x_1,x_2,x_3;q,t)\,. \tag{3.10}$$

For positive integer g the operator M^{-1} turns into the q-difference operator of order g:

$$M^{-1}:\ f(y_1,y_2,y_3)\mapsto\sum_{k=1}^{g}\,\xi_k\left(\frac{(x_1x_2)^{\frac12}}{x_3},\left(\frac{x_1}{x_2}\right)^{\frac12}\right)\,f\Big(q^{g+k}\frac{x_1}{x_3},q^{2g-k}\frac{x_2}{x_3},x_3\Big) \tag{3.11}$$

where $\xi_k(r,y)$ is given by

$$\xi_k(r,y)=(-1)^kq^{-\frac{k(k-1)}{2}}\left[\begin{smallmatrix}g\\k\end{smallmatrix}\right]_q\,y^{-2k}(1-q^{g-2k}y^{-2})\,\tfrac{(try,tr^{-1}y;q)_k(try^{-1},tr^{-1}y^{-1};q)_{g-k}}{(t^2;q)_g(q^{-k}y^{-2};q)_{g+1}}\,.$$

Remark 3. As was found in [14, 15] the operators M_ξ and M are closely related to a slightly more general integral operator $M_{\alpha\beta}$ which, in turn, is closely related to the fractional q-integration operator I^α.

Remark 4. The apparent similarity of formulas for the operator M in cases $n=2$ and $n=3$ is not a coincidence. Its explanation relies on the reduction $gl(2)\subset gl(3)$, see [15].

Acknowledgments

Authors thank M. E. H. Ismail for interesting discussions on the subject of special functions and for bringing the reference [6] to their attention. VBK wish to acknowledge the support of EPSRC.

References

[1] G. Andrews, *Summations and transformations for basic Appell series*, J. London Math. Soc. (2) **4** (1972), 618–622.

[2] R. Askey and M. E. H. Ismail, *A generalisation of ultraspherical polynomials*, in Studies in Pure Mathematics, P. Erdös, ed., Birkhäuser, 55–78, 1983.

[3] R. Askey and J. A. Wilson, Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials, Memoirs of the Amer. Math. Soc. **319**, Providence RI, 1985.

[4] M. Chakrabarty, *Formulae expressing generalised hypergeometric functions in terms of those of lower order*, Nederl. Akad. Wetensch. Proc. **A77** (1974), 199–202.

[5] R. Floreanini, L. Lapointe and L. Vinet, *A quantum algebra approach to basic multivariable special functions*, J. Phys. A: Math.Gen. **27** (1994), 6781–6797.

[6] C. Fox, *The expression of hypergeometric series in terms of similar series*, Proc. London Math. Soc. (2) **26** (1927), 201–210.

[7] G. Gasper, *Orthogonality of certain functions with respect to complex valued weights*, Cand. J. Math. **33** (1981), 1261–1270.

[8] G. Gasper and M. Rahman. Basic hypergeometric series, Cambridge University Press, 1990.

[9] R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue*, Report 94-05, Delft University of Technology, 1994.

[10] T. H. Koornwinder, *Askey-Wilson polynomials for root systems of type BC*, in: Hypergeometric functions on domains of positivity, Jack polynomials, and applications (D. St. P. Richards, ed.), Contemp. Math. **138**, AMS, Providence, R. I., (1992), 189–204.

[11] A. N. Kirillov and M. Noumi, *Affine Hecke algebras and raising operators for Macdonald polynomials*, preprint, `q-alg/9605004`.

[12] A. N. Kirillov and M. Noumi, *q-Difference raising operators for Macdonald polynomials and the integrality of transition coefficients*, preprint, `q-alg/9605005`.

[13] V. B. Kuznetsov and E. K. Sklyanin, *Separation of variables in A_2 type Jack polynomials*, RIMS Kokyuroku **919** (1995), 27–34.

[14] V. B. Kuznetsov and E. K. Sklyanin, *Separation of variables for the A_2 Ruijsenaars model and a new integral representation for the A_2 Macdonald polynomials*, J. Phys. A: Math. Gen. **29** (1996), 2779–2804.

[15] V. B. Kuznetsov and E. K. Sklyanin, *Separation of variables and integral relations for special functions*, October 1996, submitted to `The Ramanujan Journal`.

[16] L. Lapointe and L. Vinet, *Creation operators for the Macdonald and Jack polynomials*, preprint, `q-alg/9607024`.

[17] I. G. Macdonald, Symmetric functions and Hall polynomials, Oxford University Press, second edition, 1995.

[18] W. D. Niven, *On ellipsoidal harmonics*, Phil. Trans. **CLXXXII** (1891), 231.

[19] M. Noumi, *Quantum grassmannians and q-hypergeometric series*, CWI Quarterly **5** (1992), 293–307.

[20] M. Rahman and A. Verma, *Product and addition formulas for the continuous q-ultraspherical polynomials*, SIAM J. Math. Anal. 17 (1986), 1461–1474.

[21] S. N. M. Ruijsenaars, *Complete integrability of relativistic Calogero-Moser systems and elliptic function identities*, Commun. Math. Phys. **110** (1987), 191–213.

[22] E. K. Sklyanin, *Separation of variables. New trends*, Progr. Theor. Phys. Suppl. **118** (1995), 35–60.

[23] E. T. Whittaker and G. N. Watson, A course in modern analysis, Cambridge University Press, 4th ed., 1988.